

Expanded Genetic Alphabets in the Polymerase Chain Reaction**

Zunyi Yang, Fei Chen, Stephen G. Chamberlin, and Steven A. Benner*

The information content of natural DNA is limited by its construction from four nucleotide building blocks, which form just two pairs (A:T and G:C). Considerable effort has been directed towards increasing this content by developing additional nucleotides that might form additional pairs.^[1,2] One way that this can be done is by rearranging hydrogen bond donor and acceptor groups in the nucleobases to allow them to form pairs having the same geometry as A:T and G:C pairs, but that are joined by different hydrogen-bond patterns (Figure 1). This strategy can yield an artificially expanded genetic information system (AEGIS) that has as many as twelve nucleotides and as many as six base pairs.^[3]




Figure 1. Comparison of the standard C:G base pair with the non-standard Z:P pair that forms an extra two genetic letters in an artificially expanded genetic information system (AEGIS).

AEGIS nucleotides are already being applied as orthogonal binding elements in human diagnostic tools, including in FDA-approved assays for HIV, hepatitis B, and hepatitis C viruses.^[4] AEGIS oligonucleotides bind to each other and not to natural DNA, thus lowering background noise in blood and other complex biological assay mixtures.

To date, however, no practical tools have exploited additional nucleotides as components of dynamic assays in which artificial nucleotides are copied by polymerases in, for example, a polymerase chain reaction (PCR). Successful academic PCRs using expanded genetic alphabets either require a mesophilic reverse transcriptase,^[5] lose the nucleobase owing to unfavorable tautomeric equilibria,^[6] generate

highly sulfurated amplicons,^[7] or disfavor incorporation of adjacent nonstandard bases.^[8]

If it could be implemented in a fully flexible form, “six-letter PCR” should offer new approaches to analyze DNA. One challenge that has long avoided solution is multiplexed PCR, which is often complicated by off-target priming by primers presented at high concentrations in standard PCR architectures and the resulting undesired amplicons.^[9] Clever approaches to minimize such complications include various kinds of nested PCR^[10] and MegaPlex PCR.^[11] These architectures use composite primers having 3'-segments specific for their target and 5'-tags common to all primers (Figure 2a), in which PCR is initiated with composite primers in low concentration and continued using high concentrations of the common external primers that, if designed carefully, bind nothing in the assay environment. If built from A, T, G, and C, however, this design is often difficult to achieve. Essentially every 11 mer is present in the human genome, for example, meaning that essentially any sequence built from standard nucleotides will primer somewhere eventually.

Placing AEGIS nucleotides in the external tags could solve this problem. Because AEGIS-containing tags do not pair with DNA in any natural genomic background, they cannot primer to create amplicons through off-target priming, regardless of how complex the assay environment becomes. However, polymerase-AEGIS combinations are required that support fully flexible nested PCR architectures.

To develop a fully flexible nested PCR with an artificially expanded genetic alphabet, a pyrimidine analogue (6-amino-5-nitro-3-(1'- β -D-2'-deoxyribofuranosyl)-2(1H)-pyridone, **Z**) and its complementary purine analogue (2-amino-8-(1'- β -D-2'-deoxyribofuranosyl)-imidazo[1,2-a]-1,3,5-triazin-4(8H)-one, **P**; Figure 1),^[6] were examined as extra nucleotide genetic letters. Incorporated into DNA strands, these form **Z:P** pairs that contribute to duplex thermostability more than C:G pairs.^[12] Furthermore, **Z** and **P** discriminate against mismatches to the same extent as standard nucleobases,^[13] and are accepted by many polymerases.^[14]

To develop flexible nested PCR, we first addressed a concern that arises because many polymerases hold primers and templates together in their active sites sufficiently to extend the primers even when the primer–template duplex melts below the temperature used for primer extension. We deliberately designed two primers that might form dimers by **Z:G** and **P:C** mismatches (5'-CAGGAAGGAGCGATCGC-3', 5'-CAGGAAGGAGCPCATCPC-3', and 5'-CAGGAAGGAGZGATZGC-3'). We then incubated these with *Taq* polymerase to see if they generated primer dimer products (Supporting Information, Figure S1, S2). They did not, laying this concern to rest.

We then screened for polymerases that might accept PCR primers that contained multiple **Z** and **P** bases, including **Z** and **P** bases at adjacent positions. Most polymerases tested

[*] Dr. Z. Y. Yang, Dr. F. Chen, Dr. S. G. Chamberlin, Prof. Dr. S. A. Benner
Foundation for Applied Molecular Evolution
720 SW 2nd Avenue, Suite 201, Gainesville, FL 32601 (USA)
Fax: (+1) 352-271-7076
E-mail: sbenner@ffame.org
Homepage: <http://www.ffame.org>

** This work was supported by the US National Institutes of General Medical Sciences (R01M081527), the National Human Genome Research Institute (for the bioinformatics portion; R01HG004647), and the Defense Threat Reduction Agency (through its program in basic research, HDTRA1-08-1-0052).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200905173>.




Figure 2. Nested PCR with standard and AEGIS components using *Taq* DNA polymerase at four annealing temperatures (50.4, 54.2, 58.9, and 63.9°C). The initial concentration of template DNA is 0.25 nM. a) Nested PCR with different combinations of external and composite primers. Type A nested PCR: Target first amplified by standard composite primers at low concentrations, and then standard external primers (A-Std) or AEGIS external primers (A-AEGIS) at high concentrations. Type B nested PCR: Target first amplified by AEGIS composite primers at low concentrations and then standard (B-Std) or AEGIS external primers (B-AEGIS). b) Autoradiograph of a PAGE gel that resolves nested PCR products. Lanes 1–4 (A-Std): Products from composite primers and external primers, both without AEGIS. Lanes 5–8 (A-AEGIS): Absence of products with standard composite primers and AEGIS external primers. Lanes 9–12 (B-Std): Absence of products with AEGIS composite primers and standard external primers. Lanes 13–16 (B-AEGIS): Products from composite and external primers both with AEGIS.

amplified oligonucleotides containing multiple Z and P bases (Supporting Information, Figure S3). In incubation mixtures lacking dZTP and dPTP, many of these paused or stopped entirely rather than misincorporate standard triphosphates opposite Z or P; others failed to elongate efficiently following such misincorporation.

These experiments identified several polymerases that incorporated dZTP and dPTP opposite their nonstandard complements especially well and were also good at rejecting mismatches. These included Deep Vent (both exo^+ and exo^-), Vent (both exo^+ and exo^-), and *Taq* DNA polymerases (Supporting Information, Figure S3). Other polymerases, such as Phusion (a high-fidelity DNA polymerase), 9°N, Herculanase, and many *Taq* mutants also efficiently incorporate dZTP and dPTP opposite multiple template P and Z bases (data not shown).

We then investigated whether oligonucleotides containing P or Z might serve as external tags in an unplexed nested PCR using a synthetic target and *Taq* polymerases (Figure 2a). To this end, two types of experiments were designed. The first (type A) had standard nucleotides in the external tags of the composite primers (F-External-Std, R-External-Std, F-Composite-Std, and R-Composite-Std; Table 1). The second (type B) had four P bases in the external tags and composite primers replacing four G bases (F-External-4P, R-External-4P, F-Composite-4P, and R-Composite-4P; Table 1). Initial PCR amplicons obtained with these composite primers in low concentration (2.5 nM) were then further amplified with external primers in high concentration (250 nM), where the external primers were either matched to the tags used in the composition primers (the A-Std and B-AEGIS experiments) or were mismatched to the tags (the A-AEGIS and B-Std experiments). Both the A-Std (standard nested PCR, lanes 1–4) and B-AEGIS (P-containing nested PCR, lanes 13–16) experiments gave desired PCR amplicons (Figure 2b), thus showing that P-containing external and composite primers support nested PCR. In contrast, nested PCR experiments that mismatched the external primers (standard or P-containing) with the incorrect composite primers (P-containing and standard, respectively) gave no PCR products (A-AEGIS, lanes 5–8; B-Std, lanes 9–12; Figure 2b), which demonstrated the orthogonality of priming.

To further show that the P-containing external primers could not anneal and amplify standard mismatch semi-complementary sequences, we monitored the amplification in A-Std and A-AEGIS experiments using real-time PCR (Roche LightCycler 480). The A-Std experiments showed significant enhancement of fluorescence signal (SYBR green) after ten PCR cycles. In contrast, the A-AEGIS experiment gave no PCR amplicon under all annealing temperatures tested (50.4°C, 54.2°C, 58.9°C, and 63.9°C) even after 40 PCR cycles (data not shown). This clearly shows that four P:C mismatches in both forward and reverse external primers (F-External-4P and R-External-4P) prevent P-containing primers from annealing to semi-complementary C-containing sequences.

To show that nested PCR with Z- and P-containing primers can produce long amplicons, composite primers with and without P in their external tags were designed to amplify

Table 1: Oligonucleotides used in Figure 2 and Figure 3.

F-External-Std	5'-CTAGGACGACGGACTGC-3'	
F-Composite-Std	5'-CTAGGACGACGGACTGCCCATGGGAGACCGCGGT-3'	
R-Template	3'-GGTACCCTCTGGCGCCACCCGGCCGCCCATGGTAGCTATGCGCAA-5'	
F-Template	5'-CCATGGGAGACCGCGGTGGGCCCGGGTACCATCGATACGCGTT-3'	
R-Composite-Std		3'-CCATGGTAGCTATGCGCAACGCTAGCGAGGAAGGAC-5'
R-External-Std		3'-CGTAGCGAGGAAGGAC-5'
F-External-4P	5'-CTAPGACFACGFACTPC-3'	
F-Composite-4P	5'-CTAPGACFACGFACTPCCCATGGGAGACCGCGGT-3'	
R-Composite-4P		3'-CCATGGTAGCTATGCGCAACPCFAPCGAGGAAPGAC-5'
R-External-4P		3'-CPCTAPCGAGGAAPGAC-5'

the 2.5 kb gene for *Taq* DNA polymerase itself. Standard nested PCR (Figure 3, lane 3) generated the desired amplicon (2.5 kb), but with significant amounts of an undesired amplicon (750 bp, lane 4). The undesired amplicon was unassigned, but may have arisen from off-target priming involving five or six 3'-terminal matches. In contrast, nested PCR exploiting P-containing primers (lane 6) produced clean products without undesired amplicons.




Figure 3. Products from nested PCR amplification of *Taq* gene using standard or AEGIS-containing primers. Lane 1: Hi-LO™ DNA marker 50 bp–10 000 pb (Bionexus). Lane 2: Control without target; composite and external primers without AEGIS. Lane 3: Standard nested PCR with standard primers. Lane 4: Control lacking composite primers, but with standard external primers and target. Lane 5: Control without target; composite and external primers with AEGIS. Lane 6: Standard nested PCR with AEGIS primers. Lane 7: Control lacking composite primers, but with AEGIS external primers and target.

We then investigated which polymerases amplified targets most efficiently in a nested PCR architecture with composite primers and P or Z in the external tags. As a metric for evaluating real-time PCR, the polymerase generating higher fluorescence (SYBR green) at an earlier cycle was awarded a higher efficiency score, the number of cycles required to reach 50% of the end point fluorescence. The efficiency of each polymerase was further confirmed using the intensity of PCR amplicon stained by ethidium bromide on agarose gel. *Taq*, Deep Vent (both exo^+ and exo^-), Vent (both exo^+ and exo^-), Phusion, 9°N, and Herculase were tested.

For both P- and Z-containing nested PCR, Deep Vent (exo^-) was the most efficient polymerase lacking exonuclease activity (data not shown). Phusion, however, had the highest efficiency among polymerases having proofreading activity. P-containing primers gave higher nested PCR yields than Z-containing tags and primers with these polymerases (data not shown). Therefore, we further developed the nested PCR system using P-containing primers and Phusion (Supporting Information, Figure S4).




Figure 4. Seven parallel nested PCR amplifications of three genes (TOP1, HBEGF, and MYC) from human genomic DNA (Phusion high fidelity DNA polymerase) under seven different indicated annealing temperatures. Left: Amplification with standard external and composite primers. Right: Amplification using P-containing external and composite primers. Controls A (for standard nested PCR) and B (for P-containing nested PCR) were performed at 58.8°C without genomic DNA.

To demonstrate the possibility of downstream cloning of amplicons obtained with external primers containing P, an 81-nucleotide segment within the *Taq* gene was amplified with P-containing primers and Phusion. PCR products were cloned at the polylinker site of Vector pSC-B-amp/kan, and transformed (StrataClone Blunt PCR Cloning Kit) into *E. coli*. Blue-white screening gave 50 colonies that were submitted for sequencing (BioBasic, Canada); 33 sequences were returned. No point mutation errors were found in the 1377 sites sequenced, placing an upper limit on the error in this PCR (<0.1%).

PCR artifacts increase in multiplexed PCR as the number of amplicons and primers increases. We therefore tested whether nested multiplexed PCR with P-containing external primers might give cleaner products than with standard primers when applied to human genomic DNA (Promega,

G147A). Composite primers having four P bases in their external tags were synthesized to target the three cancer-associated genes examined by Li and Makrigiorgos (TOP1, HBEGF, and MYC).^[15] Sequences of gene-specific primer segments of these composites were taken from this paper (Supporting Information, Table S1).

Three-plexed nested PCR with standard primers (Figure 4, left) gave the desired amplicons, but only with substantial amounts of undesired amplicons, again presumably from off-target priming with mismatches; a similarity search suggested candidate regions for this undesired priming. At lower annealing temperatures (from 53.4 to 58.8°C), considerable primer dimer (about 40 bp) also appeared; at higher annealing temperatures (from 61.5 to 65.5°C), primer dimers disappeared, while off-target PCR artifacts did not. In contrast, three-plexed nested PCR with P-containing primers produced the desired amplicons with negligible amounts of undesired products and no primer dimers were detected under any annealing temperatures tested (Figure 4, right).

Interestingly, in controls lacking genomic DNA, 45 PCR cycles with standard primers gave primer dimers and considerable amounts of long PCR artifacts, which presumably arise from further priming from dimers (Figure 4, far left). However, the same 45 PCR cycles with P in the external and composite primers gave primer dimer (chimeric product) without further artifacts (Figure 4, far right). Furthermore, increasing the number of PCR cycles led, with standard primers, to a "PCR mess", which was not seen with P-containing external primers, even after 55 PCR cycles (Supporting Information, Figure S5).

Although we expected AEGIS to improve PCR cleanliness, we did not expect improvement to this extent. Interestingly, external tags containing P still produced better results when the analyte-specific sequences designed by Li and Makrigiorgos were replaced by external primers carefully designed by genomic analysis to bind nowhere in the human genome (Supporting Information, Table S2). Even in this case, standard external primers gave off-target priming (Supporting Information, Figure S6).

We believe that nested PCR performs better with P in the external primers because 1) P-containing external primers cannot form primer dimers with P:C mismatches, even at high concentrations; 2) P-containing external primers cannot touch down elsewhere in genomic DNA to produce off-target PCR amplicons; and 3) all the PCR amplicons end up having a P-containing sequence at their 5'-ends and a Z-containing sequence at 3'-ends; therefore, even when P-containing composite primers do produce unwanted species, they do not produce further PCR artifacts by binding to other components in the assay mixture.

In conclusion, the orthogonality of AEGIS components should improve the performance of other architectures that detect DNA in dynamic assays. Furthermore, synthetic genetic alphabets other than P and Z^[16] are now becoming available in other laboratories.^[1,8] We expect that several of these alphabets will also eventually be used to clean up multiplexed PCR. By providing the first example of a fully flexible nested PCR, however, the work described herein

should encourage their application of all artificial genetic alphabets to this goal, especially now that it is clear that AEGIS orthogonality substantially improves the quality of products produced by multiplexed PCR. This work sets the stage for the next challenges, which include demonstrating that multiple AEGIS components can be placed within the amplicon as well as in PCR tags.

Received: September 16, 2009

Published online: November 27, 2009

Keywords: base pairs · cancer · DNA replication · nucleobases · polymerase chain reaction

- [1] a) A. T. Krueger, H. G. Lu, A. H. F. Lee, E. T. Kool, *Acc. Chem. Res.* **2007**, *40*, 141–150; b) I. Hirao, *Curr. Opin. Chem. Biol.* **2006**, *10*, 622–627; c) A. A. Henry, F. E. Romesberg, *Curr. Opin. Chem. Biol.* **2003**, *7*, 727–733; d) Y. Kim, A. M. Leconte, Y. Hari, F. E. Romesberg, *Angew. Chem.* **2006**, *118*, 7973–7976; *Angew. Chem. Int. Ed.* **2006**, *45*, 7809–7812; e) H. O. Sintim, E. T. Kool, *Angew. Chem.* **2006**, *118*, 2008–2013; *Angew. Chem. Int. Ed.* **2006**, *45*, 1974–1979; f) A. M. Leconte, S. Matsuda, G. T. Hwang, F. E. Romesberg, *Angew. Chem.* **2006**, *118*, 4432–4435; *Angew. Chem. Int. Ed.* **2006**, *45*, 4326–4329; g) C. Yu, A. A. Henry, F. E. Romesberg, P. G. Schultz, *Angew. Chem.* **2002**, *114*, 3997–4000; *Angew. Chem. Int. Ed.* **2002**, *41*, 3841–3844.
- [2] S. A. Benner, *Acc. Chem. Res.* **2004**, *37*, 784–797.
- [3] a) J. A. Piccirilli, T. Krauch, S. E. Moroney, S. A. Benner, *Nature* **1990**, *343*, 33–37; b) J. D. Bain, C. Switzer, A. R. Chamberlin, S. A. Benner, *Nature* **1992**, *356*, 537–539.
- [4] a) C. A. Gleaves, J. Welle, M. Campbell, T. Elbeik, V. Ng, P. E. Taylor, K. Kuramoto, S. Accituno, E. Lewalski, B. Joppa, L. Sawyer, C. Schaper, D. McNairn, T. Quinn, *J. Clin. Virol.* **2002**, *25*, 205–216; b) T. Elbeik, J. Surtihadi, M. Destree, J. Gorlin, M. Holodniy, S. A. Jortani, K. Kuramoto, V. Ng, R. Valdes, A. Valsamakis, N. A. Terrault, *J. Clin. Microbiol.* **2004**, *42*, 563–569.
- [5] A. M. Sismour, S. Lutz, S. J.-H. Park, M. J. Lutz, P. L. Boyer, S. H. Hughes, S. A. Benner, *Nucleic Acids Res.* **2004**, *32*, 728–735.
- [6] S. C. Johnson, C. B. Sherrill, D. J. Marshall, M. J. Moser, J. R. Prudent, *Nucleic Acids Res.* **2004**, *32*, 1937–1941.
- [7] A. M. Sismour, S. A. Benner, *Nucleic Acids Res.* **2005**, *33*, 5640–5646.
- [8] a) I. Hirao, T. Mitsui, M. Kimoto, A. Yokoyama, *J. Am. Chem. Soc.* **2007**, *129*, 15549–15555; b) M. Kimoto, R. Kawai, T. Mitsui, S. Yokoyama, I. Hirao, *Nucleic Acids Res.* **2009**, *37*, e14.
- [9] J. B. Fan, M. S. Chee, K. L. Gunderson, *Nat. Rev. Genet.* **2006**, *7*, 632–644.
- [10] a) J. Brownie, S. Shawcross, J. Theaker, D. Whitcombe, R. Ferrie, C. Newton, S. Little, *Nucleic Acids Res.* **1997**, *25*, 3235–3241; b) Z. W. Lin, X. F. Cui, H. H. Li, *Proc. Natl. Acad. Sci. USA* **1996**, *93*, 2582–2587.
- [11] L. S. Meuzelaar, O. Lancaster, J. P. Pasche, G. Kopal, A. J. Brookes, *Nat. Methods* **2007**, *4*, 835–837.
- [12] Z. Y. Yang, D. Hutter, P. P. Sheng, A. M. Sismour, S. A. Benner, *Nucleic Acids Res.* **2006**, *34*, 6095–6101.
- [13] P. P. Sheng, Z. Y. Yang, Y. M. Kim, Y. R. Wu, W. H. Tan, S. A. Benner, *Chem. Commun.* **2008**, 5128–5130.
- [14] Z. Y. Yang, A. M. Sismour, P. P. Sheng, N. L. Puskar, S. A. Benner, *Nucleic Acids Res.* **2007**, *35*, 4238–4249.
- [15] J. Li, G. M. Makrigiorgos, *Nat. Protoc.* **2007**, *2*, 50–58.
- [16] Z and P are available as both triphosphates and protected phosphoramidites through reagents@ffame.org.